

A11102 129258

NAT'L INST OF STANDARDS & TECH R.I.C.




A11102129258

/Scientific papers of the Bureau of Stan
QC1 .U572 V19:1923-24 C.1 NBS-PUB-C 1919







DEPARTMENT OF COMMERCE

SCIENTIFIC PAPERS

OF THE

BUREAU OF STANDARDS

GEORGE K. BURGESS, DIRECTOR

VOLUME 19

1923-24



WASHINGTON
GOVERNMENT PRINTING OFFICE
1925

30181

QCI

- J-72

Ref-2

UNCLASSIFIED
TO
CONTRACTS
VARIABLE

CONTENTS OF VOLUME 19

	Page
469. DIRECTIVE RADIO TRANSMISSION ON A WAVE LENGTH OF 10 METERS. <i>Francis W. Dunmore and Francis H. Engel</i>	1
470. A METHOD FOR THE ACCURATE MEASUREMENT OF SHORT-TIME INTER- VALS..... <i>Harvey L. Curtis and Robert C. Duncan</i>	17
471. METHODS OF MEASUREMENT OF PROPERTIES OF ELECTRICAL INSULATING MATERIALS..... <i>J. H. Dellinger and J. L. Preston</i>	39
472. ALTERNATING-CURRENT RESISTANCE AND INDUCTANCE OF SINGLE-LAYER COILS..... <i>C. N. Hickman</i>	73
473. A METHOD FOR THE MEASUREMENT OF SOUND INTENSITY... <i>J. C. Karcher</i>	105
474. SERIES IN THE ARC SPECTRUM OF MOLYBDENUM..... <i>C. C. Kiess</i>	113
475. VISIBILITY OF RADIANT ENERGY... <i>K. S. Gibson and E. P. T. Tyndall</i>	131
476. A STUDY OF RADIO SIGNAL FADING. <i>J. H. Dellinger, L. E. Whittemore, and S. Kruse</i>	193
477. SPECTRORADIOMETRIC ANALYSIS OF RADIO SIGNALS..... <i>Chester Snow</i>	231
478. REDETERMINATION OF SECONDARY STANDARDS OF WAVE LENGTH FROM THE NEW INTERNATIONAL IRON ARC. <i>W. F. Meggers, C. C. Kiess, and Keivin Burns</i>	263
479. INTERFEROMETER MEASUREMENTS OF THE LONGER WAVES IN THE IRON ARC SPECTRUM..... <i>W. F. Meggers and C. C. Kiess</i>	273
480. A DIRECTIVE TYPE OF RADIO BEACON AND ITS APPLICATION TO NAVIGA- TION..... <i>F. H. Engel and F. W. Dunmore</i>	281
481. MEASUREMENT OF LOW RESISTANCE BY MEANS OF THE WHEATSTONE BRIDGE..... <i>Frank Wenner and Alva Smith</i>	297
482. GRAVITATIONAL ANISOTROPY IN CRYSTALS. <i>Paul R. Heyl</i>	307
483. INVESTIGATION OF THE PLATINUM METALS: IV. DETERMINATION OF IRIDIUM IN PLATINUM ALLOYS BY THE METHOD OF FUSION WITH LEAD. <i>Raleigh Gilchrist</i>	325
484. PREPARATION AND PROPERTIES OF PURE IRON ALLOYS: IV. DETERMI- NATION OF THE CRITICAL RANGES OF PURE IRON-CARBON ALLOYS BY THE THERMOELECTRIC METHOD..... <i>J. F. T. Berliner</i>	347
485. APPLICATION OF THE INTERFEROMETER TO MEASUREMENTS OF THE THER- MAL DILATATION OF CERAMIC MATERIALS..... <i>George E. Merritt</i>	357
486. SOME NEW THERMOELECTRICAL AND ACTINOELECTRICAL PROPERTIES OF MOLYBDENITE..... <i>W. W. Coblenz</i>	375
487. A QUANTITATIVE STUDY OF REGENERATION BY INDUCTIVE FEED BACK. <i>C. B. Jolliffe and Miss J. A. Rodman</i>	419
488. THERMAL EXPANSION OF MOLYBDENUM.... <i>Peter Hidnert and W. B. Gero</i>	429
489. PRIMARY RADIO-FREQUENCY STANDARDIZATION BY USE OF THE CATHODE- RAY OSCILLOGRAPH..... <i>Grace Hazen and Frieda Kenyon</i>	445
490. SPECTRA AND CRITICAL POTENTIALS OF FIFTH GROUP ELEMENTS. <i>Arthur E. Ruark, F. L. Mohler, Paul D. Foote, and R. L. Chenault</i>	463
491. THEORY OF DETERMINATION OF ULTRA-RADIO FREQUENCIES BY STAND- ING WAVES ON WIRES..... <i>August Hund</i>	487

	Page
492. FORMULAS, TABLES, AND CURVES FOR COMPUTING THE MUTUAL INDUCTANCE OF TWO COAXIAL CIRCLES... <i>Harvey L. Curtis and C. Matilda Sparks</i>	541
493. ULTRA-VIOLET REFLECTING POWER OF SOME METALS AND SULPHIDES. <i>W. W. Coblenz and C. W. Hughes</i>	577
494. ABERRATIONS OF LONG FOCUS ANASTIGMATIC PHOTOGRAPHIC OBJECTIVES <i>A. H. Bennett</i>	587
495. A RADIOMETRIC INVESTIGATION OF THE GERMICIDAL ACTION OF ULTRA-VIOLET RADIATION..... <i>W. W. Coblenz and H. R. Fulton</i>	641
496. EFFECT OF STRESS ON THE MAGNETIC PROPERTIES OF STEEL WIRE. <i>R. L. Sanford</i>	681
497. THERMAL EXPANSION OF ALUMINUM AND VARIOUS IMPORTANT ALUMINUM ALLOYS <i>Peter Hidnert</i>	697

DEPARTMENT OF COMMERCE

BUREAU OF STANDARDS

George K. Burgess, Director

SCIENTIFIC PAPERS OF THE BUREAU OF STANDARDS, No. 479

[Part of Vol. 19]

**INTERFEROMETER MEASUREMENTS OF
THE LONGER WAVES IN THE
IRON ARC SPECTRUM**

BY

W. F. MEGGERS, Physicist

C. C. KIESS, Associate Physicist

Bureau of Standards

January 5, 1924



PRICE, 5 CENTS

\$1.25 PER VOLUME ON SUBSCRIPTION

Sold only by the Superintendent of Documents, Government Printing Office
Washington, D. C.

WASHINGTON
GOVERNMENT PRINTING OFFICE

1924

INTERFEROMETER MEASUREMENTS OF THE LONGER WAVES IN THE IRON ARC SPECTRUM.

By W. F. Meggers and C. C. Kiess.

ABSTRACT.

The international system of secondary standards established by interferometer comparisons of the wave lengths of selected lines in the iron arc spectrum with the wave length of the primary standard, the red radiation from cadmium, extends, at present, from the ultra-violet to the red, but no extensive comparisons of these spectra existed for iron waves longer than 6750 Å. Using the international iron arc as a source of secondary standards and cadmium vapor lamps similar to those used in the wave-length-meter comparisons for the primary standard, new values have been obtained for 161 lines ranging in wave length from 5534.525 Å to 8824.238 Å. Seventy-five of these are longer than 6750 Å. The probable error of each value is of the order of 0.001 Å. In the region in which these determinations overlap the international standards there is a systematic deviation indicating that the accepted international scale is nearly 1 part in 1,000,000 too large. Comparison of these values with the relative ones obtained in the same spectral interval by Burns shows good agreement if the latter are adjusted to the new scale of absolute values. A figure illustrating the dispersion of phase change at reflection in interferometer mirrors of silver and copper is given.

CONTENTS.

	Page.
I. Introduction	273
II. Apparatus and methods	274
III. Results	275
IV. Discussion	280

I. INTRODUCTION.

The international system of secondary standards of wave length, as derived from the spectrum of the iron arc, now extends from the ultra-violet (3370.789 Å) to the red (6750.163 Å). During the past decade, very little progress has been made in extending this system to shorter or to longer waves. The most extensive interference measurements in the region of longer waves (5434.529 to 8824.254 Å) were made by Burns,¹ who determined the values of 125 iron lines relative to the international standards in the interval 5400 to 6500 Å. Eversheim² compared the wave lengths of 17 iron lines (6003.039 to 7445.800 Å) with the primary standard—the red radiation from cadmium. In the

¹ Jour. de Phys. (5), 3, p. 457; 1913.
63822°—24

² Ann. d. Phys., 45, p. 454; 1914.

present paper is given a table of the wave lengths of 161 lines (5434.525 to 8824.238 Å) measured by interferometer comparisons with the primary standard.

II. APPARATUS AND METHODS.

In consequence of the detection of small variations in the wave lengths of certain groups of iron lines, depending upon operating conditions, the specifications for the iron arc were recently modified by the International Astronomical Union.³

At the same time it was recommended (*loc. cit.*) that the arc previously described by the International Union for Cooperation in Solar Research⁴ be retained as a source for waves longer than 6000 Å, since the secondary standards in this region are all stable lines and exposures with the modified arc might be rather long. Accordingly, we have used an iron arc operated on a 240-volt direct-current circuit with a current of 6 amperes, the electrodes being iron rods of 7 mm diameter separated by 6 mm, and light being taken from an axial part of about 2 mm in the middle of the arc. Since, for the longer waves, the positive flame is very weak compared with that from the negative electrode⁵ the polarity was reversed during each exposure so that the positive electrode was above the negative for half the exposure and vice versa for the remaining half. This procedure gave the same photographic density to the interference fringes above and below the center of the system.

Cadmium vapor lamps of the H type similar to those used in the meter-wave-length comparisons were employed as a source of the primary standard. These lamps were made of Pyrex glass to reduce the losses due to breakage which might result from rather large changes in temperature. In actual operation, a small electric furnace held the lamp at a temperature near 300° C., and the cadmium spectrum was excited by an electrical discharge from the secondary of a transformer, whose primary was fed by 120 volts, 2 to 5 amperes alternating current.

The International Astronomical Union has adopted certain well determined values of neon wave lengths as auxiliary standards⁶ which, for practical purposes, are more convenient working standards and can be accepted as equivalent to the primary standard. Some of the exposures were made simultaneously to the iron arc

³ Trans. I. A. U., 1, p. 36; 1922.

⁴ Trans. I. U. S. R., 4, p. 58; 1914.

⁵ Fabry and Buisson, Jour. de Phys., 9, p. 229; 1910.

⁶ Trans. I. A. U., 1, p. 35; 1922.

and cadmium lamp, with the aid of a semitransparent mirror, which reflected light from the one and transmitted light from the other. In the remaining cases, in order to record the fainter lines, exposures of two hours were made to the iron arc and the primary standards were photographed on the same plate, before and after the exposure to iron. The mean value of the interferometer thickness as determined from the cadmium and neon exposures flanking that of iron was then used for reducing these plates. Three of the plates also received exposures to argon. A summary of all the observations is given in Table 1.

TABLE 1.—Summary of Observations on the Longer Waves in the Iron Arc Spectrum.

Plate.	Étalon.	Exposures of sources (in minutes).	Plate.	Étalon.	Exposures of sources (in minutes).
	mm			mm	
G1390....	5	Cd, 30; Fe, 120; A, 60; Ne, 12.	G1396....	10	Ne, 12; Fe, 120; Cd, 30; Ne, 12.
G1391....	7.5	Cd, 30; Fe, 120; A, 60; Ne, 15.	G1467....	10	Cd, Fe, simultaneously, 60.
G1392....	10	Cd, 30; Fe, 120; A, 30; Ne, 15.	G1468....	10	Cd, Fe, simultaneously, 60.
G1393....	15	Cd, 20; Ne, 15; Fe, 120; Ne, 15.	G1469....	7.5	Cd, Fe, simultaneously, 60.
G1395....	15	Cd, 45; Ne, 12; Fe, 120; Ne, 12.	G1470....	7.5	Cd, Fe, simultaneously, 60.

Schleussner ultra-rapid plates (40 cm by 6 cm) bathed in solutions of dicyanin were used for photographing the spectra. These plates were of extra thin glass, which permitted their bending to fit the best focus for lines and interference rings throughout the entire spectrum extending from about 5200 Å to 9000 Å.

The observations here described were made during 1922 when the international secondary standards (3370 to 6678 Å) were being redetermined⁷ in the same laboratory. Since the same methods and, to a large extent, the same apparatus were employed in both investigations it is unnecessary here to repeat what has been said there about the diffraction grating spectrograph, the interferometer, or the measurements and computations of wave lengths. The essential respects in which the present observations deviated from those on the shorter waves have been mentioned above so we can proceed at once to the presentation of the final results.

III. RESULTS.

All interferometer measurements of wave lengths greater than 5434 Å in the iron arc spectrum are collected in Table 2. In the first column, the estimated relative intensities of the lines are given, and where data are available the group and class to which each line belongs according to Gale and Adams⁸ is also given.

⁷ B. S. Sci. Papers, No. 478.⁸ Astroph. Jl., 35, p. 10; 1912.

Column 2 contains the new wave-length values obtained by direct comparison with the primary standard, and in the two succeeding columns the number of observations and the probable error of the mean value are given for each line. "A" indicates a probable error less than 0.0007 Å, "B," 0.0007 to 0.0012 Å, and "C" means that the determination is poor. Columns 5 and 6 present the values and probable errors published by Burns and in the next column the values obtained by Eversheim are repeated. In the last two columns are presented, for purposes of comparison with preceding values, the present values of the international secondary standards and the interpolated values adopted by the International Astronomical Union.⁹ In order to make comparisons with a considerable number of the accepted secondary standards we begin our table with 5434 Å, thus including 25 standards, 4 of which belong to group d.

TABLE 2.—Wave Lengths in the Iron Arc Spectrum.

Intensity, group, and class.	ÅB. S.	Number of observations.	Probable error.	Burns.	Probable error.	Eversheim.	Secondary standards.	Interpolated.
4 a 4.....	5434.525	6	B	529	—2	527	528
5 a 4.....	46.919	6	A	922	A	921
6 a 4.....	55.614	8	B	614	615
3 d 5.....	76.577	4	A
4 a 3.....	5497.520	10	A	520	+2	522	522
4 a 3.....	5501.469	10	A	470
5 a 3.....	06.782	10	A	784	784
2 a 4.....	35.420	2	C
2.....	65.700	2	C
6 d 5.....	69.631	10	A	633
7 d 5.....	72.856	10	B
4 d 5.....	76.102	8	A
10 d 5.....	5586.770	10	B	772
5 d 5.....	5602.959	10	B	962	C
10 d 5.....	15.658	10	B	661
5 d 5.....	24.555	10	B
2 d 5.....	38.270	3	C	276	B
5 d 5.....	58.834	8	B	836
3.....	5662.529	7	B
3.....	5701.552	8	B	553	A
4.....	09.392	10	A	395	B	396
2.....	17.845	4	C	852	B
2.....	31.770	3	C	773	B
3.....	53.138	7	B	142	A
5.....	63.009	9	A	013
2.....	5775.096	4	A	101	B
2.....	5934.675	4	B	682	D
2 d 5.....	5956.693	3	C	695	D
3.....	6003.033	7	A	036	B	039
4.....	08.577	6	B	584	B
5.....	24.060	9	B
2 b 4.....	27.056	5	C	059	059
2.....	42.088	5	C	092	B
7 b 4.....	65.489	9	A	491	492	492
3.....	78.484	3	C	485	B

⁹ Trans. I. A. U., 1, p. 41; 1922.

TABLE 2.—Wave Lengths in the Iron Arc Spectrum—Continued.

Intensity, group, and class.	λ B. S.	Number of observations.	Probable error.	Burns.	Probable error.	Ever-sheim.	Secondary stand-ards.	Inter-polated.
1.....	6089.566	4	C	570	C			
2.....	6127.910	6	B	919	B			915
5 b 4.....	36.621	10	B					624
5 b 4.....	37.699	10	B			703	701	702
2.....	51.624	4	A					
2 b 4.....	57.731	6	A	736	A			734
2.....	65.364	4	C	372	B			368
3 b 4.....	73.342	6	A	347	C			344
7 b 4.....	6191.565	10	A				568	568
3 b 4.....	6200.320	8	A	323	A			323
4 b 4.....	13.435	10	A	439	A			
4 b 4.....	19.286	10	A	289	A			290
7 b 4.....	30.730	10	A				734	734
2.....	32.661	3	C					
5 d 5.....	46.334	10	A	339	C			
5 b 4.....	52.564	10	B	567	B			567
3 b 4.....	54.262	7	B	268	B			267
3 b 4.....	56.366	7	B	372	B			
4 b 4.....	65.140	9	A	143	+2	145	145	145
2.....	80.621	6	A	625	B			
3 b 4.....	6297.800	10	B	801	A			803
5 d 5.....	6301.515	5	B					
4 b 4.....	18.025	10	A			030	028	028
2.....	22.693	6	A	697	A			696
5 b 4.....	35.338	10	A	343	-2		341	342
4 d 5.....	36.841	10	A	842	B			
2.....	44.155	6	B	158	C			
3.....	55.037	6	B	040	B			
3.....	80.748	6	A	752	C			753
7 b 4.....	6393.608	10	A				612	612
8 d 5.....	6400.018	10	B					
4.....	08.034	10	A	042	C			
6 d 5.....	11.666	10	A					
5 b 4.....	21.356	10	A					362
6 b 4.....	30.853	10	A	857	+2	856	859	859
3.....	62.732	9	A	737	A			738
2.....	75.632	6	A					639
2.....	81.878	8	A	882	B			
8 b 4.....	6494.988	10	B	991	+2	991	993	993
3.....	6518.375	6	A	378	B			382
7 b 4.....	46.247	10	A	247	A	250	252	252
3.....	75.024	6	A	032	B			029
5 b 4.....	92.922	10	A	925	B	920	928	927
4 b 4.....	6593.876	8	B					
3.....	6609.117	6	B	123	A			125
2.....	27.558	4	B					
4.....	63.447	9	A	454	A	449		455
7 b 4.....	6677.994	10	A	8.000	A	7.997	8.004	8.001
2.....	6703.573	4	A					
2.....	33.164	4	C					
4.....	50.157	10	A	164	A	163	163	165
2.....	6752.724	4	B					
2.....	6806.851	6	A					
3.....	28.612	5	A	617	C			
3.....	41.355	8	A					
3.....	43.676	8	A	681	C			
5.....	55.179	9	B	184	C			
2.....	6885.772	6	C					
3.....	6916.709	6	B	712	D			
2.....	33.628	4	C					
5.....	45.211	10	A	215	B	216		
2.....	51.271	3	B					
5.....	78.857	10	A	861	A	862		
2.....	88.531	6	A					
4.....	6999.912	5	A	932	D			

TABLE 2.—Wave Lengths in the Iron Arc Spectrum—Continued.

Intensity, group, and class.	λB. S.	Number of observations.	Probable error.	Burns.	Probable error.	Ever- sheim.	Sec- ondary stand- ards.	Inter- polated.
3.....	7022.976	7	B					
3.....	38.255	7	B	257	C			
4.....	68.418	10	B	421	A			
4.....	7090.410	10	A	416	B			
2.....	7107.464	4	A					
2.....	12.178	4	B					
5.....	30.946	10	B	958	C			
2.....	32.996	6	A					
6.....	64.472	10	B	481	B			
3.....	81.222	6	C					
8.....	7187.341	10	B	348	B	356		
7.....	7207.422	6	C	431	A	442		
2.....	19.690	6	A					
3.....	23.670	6	A	677	C			
3.....	39.896	6	A	914	D			
2.....	84.843	4	C					
3.....	88.764	6	A					
5.....	7293.073	9	B	091	C			
3.....	7307.938	5	A					
4.....	11.103	5	A					
3.....	20.694	4	C					
4.....	86.394	3	C					
6.....	7389.423	10	B	437	B			
2.....	7401.691	5	B					
7.....	11.184	10	A	192	C			
3.....	18.676	6	A					
1.....	43.031	4	B					
8.....	45.778	10	B	781	A	800		
3.....	91.678	6	B					
8.....	7495.092	10	B	106	B			
2.....	7507.300	5	C					
8.....	11.047	10	A	054	C			
4.....	31.178	10	C	192	C			
2.....	46.177	4	A					
4.....	68.931	6	A	929	D			
4.....	83.801	9	B					
6.....	7586.050	10	B					
3.....	7620.538	6	B					
2.....	53.783	5	C					
3.....	61.230	5	B					
4.....	7664.306	10	A	304	C			
3.....	7710.397	5	B					
4.....	48.282	10	A	285	C			
5.....	7780.594	8	B	597	C			
5.....	7832.233	10	B	243	C			
6.....	7937.172	10	B	182	C			
5.....	45.882	10	C	889	C			
4.....	7998.980	10	C	986	B			
1.....	8028.356	4	B					
4.....	46.084	7	B	087	B			
4.....	8085.207	4	B	219	B			
1.....	8198.960	4	C					
5.....	8220.413	10	A	422	B			
5.....	8327.069	10	A	080	B			
2.....	31.956	4	C					
4.....	8387.787	10	B	785	B			
2.....	8468.422	5	A	427	D			
1.....	8514.088	5	B					
2.....	8661.915	5	C	920	D			
3.....	8688.641	6	B	640	C			
2.....	8824.238	5	A	254				

Wave-length comparisons by means of interferometers, whose plates are covered by thin metallic films, always involve a consideration of the phenomenon known as dispersion of phase change. At reflection from metallic films light apparently penetrates the films a short distance and the amount of this penetration is some function of the wave length. When wave lengths in different spectral regions are compared it is, therefore, necessary to correct for the variation in penetration or change of phase. Methods of measuring this variation have been described by Fabry and




FIG. 1.—Dispersion of phase change at reflection from silver and copper films.

Buisson,¹⁰ who also gave examples of phase-change dispersion for metallic films of silver and nickel in the red to ultra-violet spectral interval. Extensive wave-length measurements at this bureau have involved similar investigations and the results, especially for the behaviors of different films in the infra-red, may be of interest. The method of large and small étalons¹¹ has always served us in determining the phase-change dispersion. Figure 1 illustrates the variation of the "optical surface" in interferometer films of silver and copper, the deviation of the optical surface for any particular wave length from that for the cadmium line (6438.4696 Å) being

¹⁰ Jour. de Phys. (4), 7, p. 417; 1908.

¹¹ Meggers, B. S. Bull., 12, p. 199; 1915.

plotted as ordinates and wave lengths as abscissas. The silver films were used in the redetermination of the secondary standards (see preceding paper) as well as for the longer waves reported in the present paper, and have thus been investigated throughout a wide range of spectrum (3300 to 8800 Å). The data for the copper films were obtained several years ago in wave-length measurements in the spectrum of argon.¹² They are given here because they also extended into the infra-red and show a phase-change dispersion which is strikingly different from that of silver.

IV. DISCUSSION.

In the redetermination of the secondary standards (see preceding paper) from the 12 mm iron arc it was noted that a systematic deviation, averaging 0.0049 Å, from the international values existed for the 14 lines of group b between 6027 and 6678 Å; that is, the accepted international scale appeared to be nearly 1 part in 1,000,000 too large in this interval. A similar discrepancy is revealed by the above results (column 2) from the 6 mm iron arc, the same 14 lines averaging 0.0044 Å smaller wave length than the accepted international values. The difference of 0.0005 Å between the two new series of observations may represent a real change in wave length of the so-called stable lines when the length of the arc is changed.

Since the values given by Burns were determined relative to international secondary standards, in order to make a fair comparison of his relative values with ours, the former should be reduced by about 0.005 Å at 6438 Å, and for other regions by amounts proportional to the wave-length ratios. Omitting the values which are poorly determined, such a comparison on 79 lines shows that the average difference is only ± 0.002 Å, and also that there is no appreciable systematic difference between our values and the adjusted Burns values. The agreement between our values and those of Eversheim is quite unsatisfactory, but no explanation of the deviation is offered.

WASHINGTON, August 20, 1923.

¹² Meggers, B. S. Sci. Papers, 17, p. 193; 1921.

INDEX TO VOLUME 19


	Page
A	
A quantitative study of regeneration by inductive feed back.....	419
Aberrations of photographic lenses.....	587
Absorption of radio waves.....	193
spectra. <i>See</i> Spectra.	
Actinoelectrical properties of molybdenite ...	375
Alternating current resistance and inductance of solenoids.....	73
current theory of regeneration.....	419
Alloy, iron-carbon, determination of critical ranges by thermoelectric means.....	347
Alloys, determination of iridium in.....	325
platinum.....	325
preparation of.....	325
Aluminum alloys, thermal expansion.....	697
copper alloys, thermal expansion.....	697
manganese alloys, thermal expansion.....	697
copper alloys, thermal expansion.....	697
silicon alloys, thermal expansion.....	697
copper alloys, thermal expansion.....	697
manganese alloys, thermal expansion.....	697
thermal expansion.....	697
zinc alloys, thermal expansion.....	697
Amateur, effect of fading on reception by....	193
radio wave finding.....	193
Antenna, airplane, directive receiving properties of.....	281
double coil.....	281
double coil, directional transmission, characteristics of.....	281
loop, transmitting.....	281
Antimony, spectral classifications, critical potentials, and absorption spectra.....	463
Applications of the interferometer.....	357
Arc spectrum of molybdenum.....	113
Arsenic, spectral classifications, critical potentials, and absorption spectra.....	463
Atmospheric electricity, relation to radio transmission.....	193
Atmospherics, relation to radio reception....	193
B	
Barometric conditions, effect on radio transmission.....	193
Beacon, radio, directive.....	281
<i>Bennett, A. H.</i> , Aberrations of long focus anastigmatic photographic objectives.....	587
<i>Berliner, J. F. T.</i> , Preparation and properties of pure iron alloys: IV. Determination of the critical ranges of pure iron-carbon alloys by the thermoelectric method.....	347
Bismuth, spectral classifications, critical potentials, and absorption spectra.....	463
Brinell hardness numerals, table of.....	39

	Page
Calculation of inductance	641
Cathode-ray oscillograph	445
use in frequency standardization	445
<i>Chenault, R. L., Arthur E. Ruark, F. L. Mohler, Paul D. Foote and, Spectra and</i> <i>motional potentials of five group elements...</i>	463
Circuits, radio-frequency, for measurements ..	39
Coaxial circles, curves for estimating the mutual inductance	541
formulas determining mutual inductance ..	541
mutual inductance of	541
tables for computation of mutual inductance	541
<i>Coblentz, W. W., Some new thermoelectric</i> <i>and actinoelectric properties of molybdenite</i>	375
—, and <i>H. R. Fulton, A radiometric investigation</i> <i>of the germicidal action of ultra-violet</i> <i>radiation</i>	641
—, and <i>C. W. Hughes, Ultra-violet reflecting</i> <i>power of some metals and sulphides</i>	577
Coil antenna, transmitting	281
Coils, single-layer, alternating-current, resistance and inductance of	73
Ceramics	357
Critical potentials of arsenic, antimony, and bismuth ranges in iron-carbon alloys, determined by thermoelectric means ...	347
Crystals	307
<i>Curtis, Harvey L., and C. Matilda Sparks,</i> <i>Formulas, tables, and curves for computing</i> <i>the mutual inductance of two coaxial circles</i>	541
—, and <i>Robert C. Duncan, A method for the</i> <i>accurate measurement of short-time intervals</i>	17
Curves	541

D	
"Dead spots" and radio reception.....	193
Decrement	231
<i>Dellinger, J. H., and J. L. Preston, Methods of measurement of properties of electrical insulating materials.</i>	39
—, <i>L. E. Whittemore, and S. Kruse, A study of radio signal fading.</i>	193
Density, electrical insulating materials, measurement.....	39
Dielectric constant, insulating materials, measurement.....	39
Directional variations of radio signals.....	193
Directive radio transmission.....	281
measurements.....	1
Distortion, radio wave.....	281
Diurnal variations of radio signals.....	193
<i>Duncan, Robert C., Harvey L. Curtis and, A method for the accurate measurement of short-time intervals.</i>	17

	Page		Page
<i>Dunmore, F. W., F. H. Engeland</i> , A directive type of radio beacon and its application to navigation.....	281	Heavieside layer, influence on radio transmission.....	193
—, and <i>Francis H. Engel</i> , Directive radio transmission on a wave length of 10 meters..	1	<i>Heyl, Paul R.</i> , Gravitational anisotropy in crystals.....	307
Duralumin, thermal expansion.....	697	<i>Hickman, C. N.</i> , Alternating-current resistance of single-layer coils.....	73
ultra-violet reflecting power.....	577	<i>Hidnert, Peter</i> , Thermal expansion of aluminum and various important aluminum alloys.....	697
		—, and <i>W. B. Gero</i> , Thermal expansion of molybdenum.....	429
E		High-frequency resistance, measurement.....	39
Electrical properties of insulating materials, measurement.....	39	<i>Hughes, C. W., W. W. Coblentz and</i> , Ultra-violet reflecting power of some metals and sulphides.....	577
Einstein.....	307	Humidity control tank.....	39
Energy distribution.....	231	<i>Hund, August</i> , Theory of determination of ultra-radio frequencies by standing waves on wires.....	487
<i>Engel, Francis H., Francis W. Dunmore and</i> , Directive radio transmission on a wave length of 10 meters.....	1		
—, and <i>F. W. Dunmore</i> , A directive type of radio beacon and its application to navigation.....	281	I	
Expansion, thermal, aluminum.....	697	Impact strength, electrical insulating materials, measurement.....	39
molybdenum.....	429	Inductance, alternating current, of solenoids..	73
Eye, sensibility.....	131	mutual.....	541
		Inductive coupling.....	419
F		feed back, study of.....	419
Fading, effect on radio reception.....	193	Insulating materials, electrical, properties, measurement.....	39
theory of.....	193	Interference.....	231
Flash-over voltage, radio-frequency measurement.....	39	Intensity measurements, sound.....	105
Fog signals, radio.....	281	Inverse fading of radio signals.....	193
<i>Foote, Paul D., Arthur E. Ruark, F. L. Mohler, R. L. Chenault and</i> , Spectra and critical potentials of fifth group elements..	463	Iridium.....	325
Formulas.....	541	determination in platinum alloys.....	325
Four-terminal resistors.....	297	spectrographic examination of.....	325
Frequency standardization by means of parallel wires.....	487	Iron.....	325
<i>Fulton, H. R., W. W. Coblentz and</i> , A radio-metric investigation of the germicidal action of ultra-violet radiation.....	641	arc spectra.....	263
		spectrum.....	273
G		carbon alloys, thermal analysis of thermoelectric method of thermal analysis..	347
Galena, ultra-violet reflecting power.....	577	separation from iridium.....	325
<i>Gero, W. B., Peter Hidnert and</i> , Thermal expansion of molybdenum.....	429		
Generation of very high frequency currents..	1	J	
Generator for ultra-radio frequency currents..	1	<i>Jolliffe, C. B., and Miss J. A. Rodman</i> , A quantitative study of regeneration by inductive feed back.....	419
<i>Gibson, K. S., and E. P. T. Tyndall</i> , Visibility of radiant energy.....	131		
<i>Gilchrist, Raleigh</i> , Investigations on the platinum metals: IV. Determination of iridium in platinum alloys by the method of fusion with lead.....	325	K	
Glasspots, thermal expansivities of.....	357	<i>Karcher, J. C.</i> , A method for the measurement of sound intensity.....	105
Gold, effect on determination of iridium....	325	<i>Kevin, Burns, W. F. Meggers, C. C. Kiess and</i> , Determination of secondary standards of wave length from the new international iron arc.....	263
Gravitation.....	307	<i>Kenyon, Frieda, Grace Hazen and</i> , Primary radio-frequency standardization by use of the cathode-ray oscillograph.....	445
Graphite, ultra-violet reflecting power.....	577	<i>Kiess, C. C.</i> , Series in the arc spectrum of molybdenum.....	113
		—, <i>W. F. Meggers, Kevin Burns and</i> , Determination of secondary standards of wave length from the new international iron arc..	263
H		—, <i>W. F. Meggers and</i> , Interferometer measurements of the longer waves in the iron arc spectrum.....	273
Hardness, electrical insulating materials, measurement.....	39	<i>Kruse, S., J. H. Dellinger, L. E. Whittemore and</i> , A study of radio signal lading.....	193
Hartmann test for lenses.....	587		
<i>Hazen, Grace, and Frieda Kenyon</i> , Primary radio-frequency standardization by use of the cathode-ray oscillograph.....	445		

L		R	
	Page		Page
Lenses, monaxial aberrations.....	587	Radiant energy, visibility of.....	131
Low-resistance measurements.....	297	Radio.....	231
Luminous efficiency of radiant energy.....	131	beacon.....	281
M		fading.....	193
Magnetic, properties and mechanical stress..	681	frequency properties, insulating ma-	
Measurement of resistance.....	297	terials, measurement.....	39
properties electrical insulating materials.	39	standardization.....	445
Measurements of thermal dilatations.....	357	transmitting set.....	281
Mechanical properties, electrical insulating		Reception, radio, on airplanes.....	281
materials, measurement.....	39	Reflecting power, pyrites, stibnite, molybde-	
<i>Meggers, W. F., C. C. Kiess, and Kevin</i>		nite, galena, graphite, duralumin.....	577
<i>Burns</i> , Determination of secondary stand-		Reflection of radio waves.....	193
ards of wave length from the new inter-		of very short electric waves.....	1
national iron arc.....	263	Refraction of radio waves.....	193
—, and <i>C. C. Kiess</i> , Interferometer measure-		Regeneration.....	419
ments of the longer waves in the iron arc		Resistance, alternating-current, of solenoids..	73
spectrum.....	273	measurements.....	297
<i>Merritt, George E.</i> , Application of the inter-		radio-frequency, measurement.....	39
ferometer to measurements of the thermal		variation method, radio measurements..	39
dilatation of ceramic materials.....	357	Resistivity, insulating materials, measure-	
Meteorological conditions, effect on radio		ment.....	39
transmission.....	193	Rhodium, effect on determination of iridium.	325
Microstructure, molybdenum.....	429	<i>Rodman, Miss J. A., C. B. Jolliffe and, A</i>	
<i>Mohler, F. L., Arthur E. Ruark, Paul D.</i>		quantitative study of regeneration by induc-	
<i>Foote, R. L. Chenault, and</i> , Spectra and		tive feed back.....	419
critical potentials of fifth group elements...	463	<i>Ruark, Arthur, F. L. Mohler, Paul D. Foote,</i>	
Moisture absorption, electrical insulating ma-		and <i>R. L. Chenault</i> , Spectra and critical	
terials, measurements.....	39	potentials of fifth group elements.....	463
Molybdenite, thermoelectrical and actino-		Ruthenium, effect on determination of	
electrical properties.....	375	iridium.....	325
ultra-violet reflecting power.....	577	S	
Molybdenum, series in the arc spectrum of..	113	Safety, method of improvement in aerial and	
thermal expansion.....	429	marine navigation.....	281
Mutual inductance calculations.....	641	<i>Sanford, R. L.</i> , Effect of stress on the magnetic	
of circular circuits.....	641	properties of steel wire.....	681
of circular filaments.....	641	Screen, shielding, radio measurements.....	39
of coaxial circles.....	641	Seasonal variation of radio signals.....	193
determinations.....	541	Sensibility of eye.....	131
N		Series in molybdenum.....	113
Navigation, aerial, radio, aid to.....	281	Shielding, radio measuring circuits.....	39
marine, radio, aid to.....	281	Short-time intervals.....	17
Nitrogen, spectral classifications.....	463	Short-wave directive transmission.....	1
Nocturnal variation of radio signals.....	193	Short waves, susceptibility of fading.....	193
O		Signal, radio, effect of fading.....	193
Oscillograph, cathode-ray.....	445	strength, variation of.....	193
P		<i>Smith, Alva, Frank Wenner and</i> , Measure-	
Palladium, effect on determination of iri-		ment of low resistance by means of the	
dium.....	325	Wheatstone bridge.....	297
Parabolic reflector for directive transmission.	1	<i>Snow, Chester</i> , Spectroradiometric analysis	
Parallel wire system.....	487	of radio signals.....	231
Photographic lenses, aberrations.....	587	Solenoids, alternating-current, resistance and	
Power loss, electrical insulating materials,		inductance of.....	73
measurement.....	39	Sound intensity measurements.....	105
Poynting clamp.....	307	<i>Sparks, C. Matilda, Harvey L. Curtis and</i> ,	
<i>Preston, J. L., J. H. Dellinger and</i> , Methods		Formulas, tables, and curves for computing	
of measurement of properties of electrical		the mutual inductance of two coaxial	
insulating materials.....	39	circles.....	541
Primary radio-frequency standardization...	445	Specimens, electrical insulating materials,	
by use of the cathode-ray oscillo-		preparation.....	39
graph.....	445	Spectra, spectral classifications, and excita-	
standard wave meters.....	445	tion of spectra of arsenic, antimony, bis-	
wave meter standardization.....	445	muth, and nitrogen.....	463
Progressive fading of radio signals.....	193	Spectrum analysis.....	231
Pyrites, ultra-violet reflecting power.....	577	visibility of radiant energy in.....	131



Mar 1871





